

## Opinion

## Monotopic Membrane Proteins Join the Fold

Karen N. Allen <sup>1,2,\*</sup> Sonya Entova <sup>3</sup> Leah C. Ray <sup>2</sup> and Barbara Imperiali <sup>3,4,\*</sup>

**Monotopic membrane proteins, classified by topology, are proteins that embed into a single face of the membrane. These proteins are generally underrepresented in the Protein Data Bank (PDB), but the past decade of research has revealed new examples that allow the description of generalizable features. This Opinion article summarizes shared characteristics including oligomerization states, modes of membrane association, mechanisms of interaction with hydrophobic or amphiphilic substrates, and homology to soluble folds. We also discuss how associations of monotopic enzymes in pathways can be used to promote substrate specificity and product composition. These examples highlight the challenges in structure determination specific to this class of proteins, but also the promise of new understanding from future study of these proteins that reside at the interface.**

## Introduction

**Integral membrane proteins** (see [Glossary](#)) are implicated in a multitude of essential biological processes including metabolism, biosynthesis, and cellular signaling [1]. They constitute between 20% and 30% of all proteomes [2,3]. Membrane proteins can be classified based on topology, which describes the physical arrangement of the structures with respect to the membrane (see [Figure 1A](#) in [Box 1](#)). There are three topologically distinct classes of integral membrane proteins – **polytopic**, **bitopic**, and **monotopic** [4] – all of which are irreversibly associated with the membrane and require addition of detergents for solubilization [5,6]. Membrane proteins may adopt bitopic or polytopic geometries to satisfy functional properties. For example, cell-surface receptors such as receptor tyrosine kinases and G protein-coupled receptors transduce information from the outside to the inside of cells. Similarly, channels and transporters move materials across membranes to perform critical tasks. By contrast, the relationships between topology and function are less clearcut for monotopic membrane proteins, and they are the least well understood from a structural perspective. Additionally, some proteins, categorized as monotopic membrane proteins, may be only peripherally associated thus adding to ambiguities that can be resolved only with detailed experimental analysis [7,8]. Only a scant ~0.06% of nonredundant structures exhibit the monotopic membrane topology, although integral membrane proteins as a whole constitute 4.6% (see [Figure 1A](#) in [Box 1](#)). The structures of full-length monotopic membrane proteins have proved to be more recalcitrant to structure determination, and computational methods to enable distinction between bitopic and monotopic membrane proteins are also needed.

## Location, Location, Location

The structurally characterized monotopic membrane proteins defined to date specialize in catalyzing reactions on membrane-resident substrates, often including a second hydrophilic substrate. Thus, information on the enzyme substrates and reaction mechanisms provides a valuable starting point for understanding how the structures are purposed for catalysis. There are two principle strategies for the interaction of monotopic enzymes with their substrates: enzymes may ‘extract’ hydrophobic substrates from the lipid bilayer for catalysis at an active

## Highlights

Monotopic proteins are underrepresented in the PDB, with only 25 nonredundant structures currently constituting ~0.06% of known structures.

Many monotopic membrane proteins are homologous to soluble counterparts and use common structural features to embed shallowly in the membrane.

Selected monotopic proteins engage more deeply in the membrane (e.g., associating via reentrant-helical domains).

Monotopic enzymes are purposed for catalysis of reactions involving hydrophobic or amphiphilic substrates not readily soluble in water.

The active sites of monotopic enzymes may be at the membrane surface or distal to it, and the requirement for hydrophobic substrate extraction is dictated by the substrate and the relative orientations of the active site and the membrane.

Association of multiple monotopic enzymes in pathways can be advantageously applied in the assembly of complex glycoconjugates.

<sup>1</sup>Department of Chemistry, Boston University, Boston, MA 02215, USA

<sup>2</sup>Program in Biomolecular Pharmacology, Boston University School of Medicine, Boston, MA 02118, USA

<sup>3</sup>Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

<sup>4</sup>Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

\*Correspondence: [drkallen@bu.edu](mailto:drkallen@bu.edu) (K.N. Allen) and [imper@mit.edu](mailto:imper@mit.edu) (B. Imperiali).

site positioned distal to the bilayer interface or they adopt an energetically favorable ‘leave-in-place’ strategy whereby hydrophobic and amphiphilic substrates are not subject to translocation from the membrane for catalysis (see Figure 1B in Box 1) [1]. Irrespective of the mode of substrate interaction, examination of monotopic proteins of known structure reveals common themes for membrane association (see Figure 1C in Box 1 and Figure 1). Membrane association may occur through hydrophobic interactions supported most commonly through **amphipathic** helices positioned parallel to the membrane plane [9]. Additional structural features enlisted include hydrophobic loops extending into one leaflet of the membrane and hydrophobic patches coupled with electrostatic interactions with phospholipid head groups [6,10–13]. Alternatively, the protein structures may penetrate more deeply into the hydrophobic core of the membrane, for example, through reentrant motifs that associate with both the membrane core and domains of the protein that are positioned at the membrane interface [14,15]. Molecular dynamics simulations have shown that the membrane associations of monotopic proteins that extend deeper into the membrane result in local perturbations in the lipid bilayer [10], which can be functionally important.

In addition to these specific membrane-association modalities, the known structures of monotopic membrane proteins may share some global features (see Figure 1C in Box 1). They may comprise extensive soluble domains with homology to known soluble proteins, and minimal membrane-embedded domains that have been repurposed for catalysis at the membrane interface (Figure 2 and Table 1). Many are biologically active as dimers [16] and some feature long hydrophobic tunnels connecting the membrane interface to active sites located at a distance from the membrane, allowing access to hydrophobic substrates. The majority of monotopic membrane enzymes that have been structurally characterized to date share homology with existing soluble folds of known function [17,18]. In some cases, the soluble and membrane-associated forms of enzymes are even **isoforms** of each other [19,20]. By contrast, the structures of some monotopic membrane proteins are unique, indicating that they may have evolved via membrane-dependent processes and reveal new folds not observed in soluble **homologs** [21,22].

### Double Feature

As first noted for fatty acid amide hydrolase (FAAH) [6,23], membrane association in monotopic membrane proteins is commonly enabled via insertion of membrane-interacting segments and/or dimerization to increase the surface area of the protein–membrane interaction. This strategy is also exemplified by RPE65, an enzyme expressed in the retinal pigment epithelium that catalyzes the conversion of all-*trans*-retinyl esters to 11-*cis*-retinol (Figure 3A). An especially ‘physiologically relevant’ crystal form was obtained by crystallizing RPE65 at high lipid:detergent ratios, resulting in RPE65 that is not separated from native phospholipids [24]. Thus, the isomerase packs as a dimer with the membrane-binding surface facing micelle sheets, mimicking the assembly at the membrane. Sequence- and structure-based alignments show that a ~25-amino-acid insert in RPE65 that affords the dimerization domain is not found in soluble, monomeric homologs (e.g., apocarotenoid oxygenase) [25]. Notably, **orthologs** of RPE65 from bacteria can bind membrane through a hydrophobic patch, but are monomeric and do not contain the dimer-mediating sequence. Thus, dimerization is not a requisite for the evolution of membrane binding and RPE65 and vertebrate family members may have elaborated a more extensive membrane-binding surface in concert with acquiring a dimer interface.

Another requirement of the dimerization strategy is met by FAAH, where a 180° rotation in subunit orientations within the dimer, compared with soluble dimeric family members, allows simultaneous access of both active sites to the membrane interface. Interface remodeling is not

### Glossary

**Amphipathic:** a molecule, protein segment, or protein having both hydrophobic and hydrophilic components.

**Bitopic protein:** a membrane protein that spans the membrane bilayer a single time; sometimes referred to as a ‘single-pass’ transmembrane protein.

**Glycosyl transferase (GTase):** an enzyme that catalyzes the transfer of a sugar from an activated sugar donor, commonly a nucleoside diphosphate derivative, to an acceptor such as another sugar or a lipid.

**Homologs:** members of the same protein family.

**Integral membrane protein:** a protein that is inserted into the membrane and can be solubilized only by detergent.

**Isoforms:** two structurally and functionally similar proteins with nonidentical sequences due to encoding by different genes or splicing variations.

**Monotopic protein:** a membrane protein that does not span the membrane bilayer, but rather enters and exits on a single face of the membrane bilayer.

**Orientations of Proteins in Membranes (OPM) database:** a curated web resource providing information on the position of proteins in membranes. Included in the database is output from the PPM server.

**Orthologs:** members of the same protein family having the same function but occurring in different organisms.

**Paralogs:** members of the same protein family having different functions.

**Peripheral membrane protein:** a protein associated with the membrane surface in a manner that can be dissociated by the addition of high salt.

**Phosphoglycosyl transferase (PGT):** an enzyme that catalyzes the transfer of a phosphosugar from an activated sugar donor, commonly a nucleoside diphosphate derivative, to an acceptor substrate such as a polyprenol phosphate.

**Polyprenol:** a linear long-chain alcohol featuring greater than six isoprene units (branched five-carbon units) of either *cis* or *trans* configuration.

always needed, as is the case for cyclooxygenase-2 (COX-2), also called prostaglandin H<sub>2</sub> synthase (PGHS) [26], which catalyzes the first committed step of prostaglandin biosynthesis, converting arachidonic acid to prostaglandin H<sub>2</sub> (Figure 3B). The catalytic domain of COX-2 has low sequence identity (~20%) but high structural similarity (RMSD = 2.6 Å) to the soluble mammalian peroxidases (e.g., canine myeloperoxidase) [27]. The subunit interface of the COX-2 dimer allows the same relative orientations of the catalytic domains as the subunit interface in the soluble mammalian peroxidases, with sequence inserts affording membrane interaction. Thus, nature has leveraged existing monomeric and dimeric soluble domains to allow direct access of these catalytic domains to the membrane interface, allowing extraction of substrate and release of product to the bilayer.

COX-2 [26] and  $\alpha$ -dioxygenase ( $\alpha$ -DOX) (Figure 3C) [28] illustrate a case where two enzymes from the same superfamily adopt distinct mechanisms of membrane interaction.  $\alpha$ -DOX converts linoleic acid (LA18:2  $\omega$ -6,9) and related fatty acids into 2(*R*)-hydroperoxides, which undergo spontaneous decarboxylation to chain-shortened aldehydes [29] (Figure 3C).  $\alpha$ -DOX has a two-domain structure with a membrane-interacting domain and a catalytic domain. There is significant structural similarity in the catalytic domains of  $\alpha$ -DOX and COX-2 (RMSD = 1.7 Å). Four  $\alpha$ -helices constitute a hydrophobic fatty-acid binding channel in the catalytic domain of  $\alpha$ -DOX, which spans ~20 Å between the membrane domain opening and the catalytic site. The structural arrangement of the four helices and hence the fatty-acid-binding channel is similar to that of the substrate-binding channel of COX-2. By contrast, there is no sequence or structural similarity between the membrane-binding domains of the two folds, although both interact with the membrane via amphipathic helices. These  $\alpha$ -helices tether  $\alpha$ -DOX and COX-2 to one leaflet of the membrane and serve as the access point for entry of the substrate into the catalytic domain. Together, these findings suggest that members of the cyclooxygenase–peroxidase family employ a diversity of modalities for membrane interaction. Why COX-2 forms a dimer whereas  $\alpha$ -DOX exists as a monomer is unclear from the standpoint of energetics, as the two interfaces bury similar surface areas in the membrane.

### A Difficult Extraction

The localization of monotopic proteins at the membrane interface positions them in proximity to lipid-soluble substrates, yet it is not uncommon for these enzymes to extract their substrates through hydrophobic channels into polar active sites distal to the interface for catalysis. Although energetically unfavorable, this strategy is favorable in terms of evolution, leveraging the recruitment of a soluble domain already optimized to enact catalysis proficiently. The nature of the connection between the catalytic site and the membrane interface is critical in obviating the need to transfer substrate through aqueous cellular compartments and thus has conserved features among monotopic membrane proteins that extract their membrane-soluble substrates. In RPE65, the side chains of residues lining a channel from the active site to the membrane form a continuous aromatic surface that, by interacting with the polyene chain of retinal ester, allows the uptake of substrate directly from the membrane (a distance of ~20 Å). In addition to the presence of a hydrophobic substrate channel, the features of the interface between the channel entrance and the membrane-binding surface are similar in the monotopic membrane proteins FAAH, squalene-hopene cyclase (SHC) [30] (Figure 3D), and COX-2/ $\alpha$ -DOX, despite all three examples originating from different fold families. These enzymes have basic amino acids surrounding the substrate channel entrance, allowing the binding of negatively charged phospholipid head groups constituting the membrane. Monotopic folds with hydrophobic channels to the active site have the added advantage of protection from bulk solvent. For instance, the terpene cyclases SHC and oxidosqualene cyclase contain highly conserved hydrophobic active sites containing electron-rich amino acids that allow the

**Polytopic protein:** a membrane protein that spans the membrane bilayer multiple times.

**PPM server:** calculates rotational and translational positions of transmembrane and peripheral proteins in the membrane using the PDB coordinate files as input.


**Reentrant membrane helix:** a discontinuous hydrophobic helix that enters and exits on a single face of the membrane bilayer.

**Rossmann fold:** a fold characterized by repeating motifs of  $\alpha$ -helix– $\beta$ -strand– $\alpha$ -helix secondary structure elements where the  $\beta$ -strands together form a parallel  $\beta$ -sheet.

**TMHMM:** an algorithm to predict transmembrane helices in proteins using a hidden Markov model.

## Box 1. Classification, Characteristics, and Structural Representation of Membrane Proteins

Integral membrane proteins are classified as polytopic, bitopic, or monotopic dependent on their membrane-associated topology and domain distribution across the membrane. **Peripheral proteins** associate reversibly with the lipid bilayer (Figure 1A, left). The distribution of membrane protein structures in the PDB shows a paucity of monotopic membrane protein structures (Figure 1A, right). Numbers for each class are compiled from nonredundant (70% sequence ID) PDB, **Orientations of Proteins in Membranes (OPM)**, and MPStruct databases: soluble proteins – 34 987; peripheral proteins – 1350; integral proteins – 1679; polytopic – 1399; bitopic – 256; monotopic – 25. For a curated database of membrane protein structures, see <http://blanco.biomol.uci.edu/mpstruc/>. The modes of interaction of monotopic enzymes with membrane and substrates are diagrammed in Figure 1B. FAAH exemplifies enzymes that extract the membrane-soluble substrate into the active site above the membrane interface. PglC exemplifies enzymes where the active site is at the interface, obviating extraction. In enzymes like PglH, active sites are distal to the membrane surface, allowing the correct spacing for membrane-resident substrates with hydrophilic moieties. WaaA undergoes a conformational change bringing the active site to the membrane interface. Comparison of the monotopic membrane proteins highlights shared features including oligomerization states, modes of interaction with membrane and substrate(s), and homology to soluble folds (Figure 1C).




## Comparison of features of monotopic membrane proteins

Characteristic	Monotopic enzyme
Functional dimer	NAPE-PLD, COX-2, AOX, FAAH, GlpD, SHC, RPE65, NDH-2, TagF
Amphipathic helices	CPT-II, COX-2, AOX, CYP19A1, $\alpha$ -DOX, FAAH, ETF-QO, PlsC, PglC, SHC, PglH, SQR, LpxK, NDH-2, TagF
Long hydrophobic loop	NAPE-PLD, RPE65, NDH-2
Hydrophobic 'channel' to active site	CPT-II, COX-2, CYP19A1, $\alpha$ -DOX, FAAH, SHC, GlpD, RPE65
Hydrophobic patch surrounded by positive charges	COX-2, $\alpha$ -DOX, FAAH, GlpD, WaaA, SHC
Reentrant membrane helix	LpxM, PlsC, PglC
Homology to soluble folds	All except AOX and PglC

<sup>a</sup>Abbreviations, Protein Data Bank accession code, and references as in Table 1.

Figure 1. Occurrence, Function, and Features of Monotopic Membrane Proteins.




## Trends in Biochemical Sciences

**Figure 1. Structures of Representative Monotopic Membrane Enzymes.** Protein surface colored by normalized consensus hydrophobicity from hydrophobic (red) to hydrophilic (white). Membrane represented by gray box. Location of membrane plane calculated with PPM server (<http://opm.phar.umich.edu/server.php>) for 1CQE (COX-2), 2SQC (SHC), 2QCU (GlpD), 2H4T (CPT-II), 5KYM (PlsC), 2XCI (WaaA), 5W7L (PglC), and 3EQM (CYP19A1).

generation, propagation, and shielding of carbocation intermediates from bulk solvent, preventing termination of the cyclization cascades by nucleophilic addition of water [30–32]. Similarly, the active sites and heme prosthetic groups of the cytochrome P450s are buried (e.g., at a distance of 28 Å from the membrane interface to the heme iron in CYP19A1 [33]). This feature prevents decomposition of the iron–oxygen intermediates by uncoupling reactions that consume NAD(P)H and result in the formation of reactive oxygen species [34].

Membrane-bound cytochrome P450s are targeted to the interface by an N-terminal leader sequence that includes an amphipathic helix linked by a polar connector to the catalytic domain, which is structurally homologous to that of soluble cytochrome P450s [35]. Compared with these soluble homologs, in membrane-bound cytochrome P450s (except the CYP51 family) there is a shift in the location of the beta-domain and the A-propionate heme side chain towards the proximal side of the heme – a design feature that allows access of substrates from the membrane interior [34]. To control reactivity, cytochrome P450s are structurally dynamic, with open and closed conformers for substrate association and product dissociation. Molecular dynamics studies support a model where interaction with the membrane promotes the open state of the substrate tunnel leading from the membrane interior to the buried active site [36]. In substrate-specific cytochrome P450s, lipophilic-substrate binding alters the protein dynamics and increases catalytic specificity and efficiency. Additionally, consequences of the membrane-interaction mode on conformation were revealed in the structure of a full-length fungal cytochrome P450, which contains an extended N-terminal transmembrane helix (designated as bitopic) [35]. The structure shows a rigid interaction between the membrane-resident domain



## Trends in Biochemical Sciences

**Figure 2. Comparison of Membrane Proteins and Their Soluble Homologs.** Ribbon diagram of membrane proteins (purple) superimposed with the fold core of a prototypical soluble homolog (gray). Red broken line indicates approximate membrane location calculated using the PPM server [49]. (A) Lysophosphatidic acid acyltransferase, PlsC (purple, 5KYM) and glycerol-3-phosphate (1)-acyltransferase (gray, 1K30); (B) tetraacyldisaccharide-1-phosphate 4'-kinase, LpxK (purple, 4EHW) and GTP-binding protein HypB (gray, 2HF9); (C) fatty acid alpha-deoxygenase,  $\alpha$ -DOX (purple, 4HHS), and myeloperoxidase (gray, 1CXP) with heme group in pink space filling.

and the catalytic domain, suggesting that the transmembrane domain itself may promote the catalytically competent pose [37].

As a final example, *sn*-glycerol-3-phosphate dehydrogenase (GlpD) is an essential FAD-dependent enzyme that catalyzes the oxidation of glycerol 3-phosphate (G-3-P) to dihydroxyacetone phosphate (DHAP) (Figure 3E). The enzyme is a dimer comprising a soluble extramembranous C-terminal domain and an N-terminal FAD-binding domain that includes the membrane-resident region. GlpD similarly exemplifies both the adaptation of catalytic units to act at the membrane and the extraction of a hydrophobic substrate from the membrane by both subunits of a dimer. Notably, GlpD can exist as a soluble isoform and a membrane-bound form with sequence identity >40% [19]. Based on the distribution of polar and hydrophobic residues, it is estimated that GlpD embeds to depths of 12–15 Å into the lipid bilayer and the distance from the membrane surface to the isoalloxazine ring of FAD is ~18 Å. As in many of the dimeric monotopics, the molecular two-fold axis relating the subunits within the dimer is perpendicular to the membrane, allowing access from the membrane to both active sites.


Table 1. Monotopic Membrane Proteins with Corresponding PDB Coordinate Accession Code (PDB ID) of a Representative Ortholog, Accession Code of a Soluble Homolog, and Oligomeric State

Monotopic	PDB ID	Soluble homolog	Oligomeric state
<i>N</i> -Acylphosphatidylethanolamine-hydrolyzing phospholipase D (NAPE-PLD)	4QN9 [58]	1BMC [59]	Dimer
Carnitine-palmitoyl transferase (CPT-II)	2H4T [60]	1NDB [61]	Monomer
COX-2/prostaglandin H2 synthase (COX-2/PGHS)	1CQE [26]	1CXP [62]	Dimer
Cyanide-insensitive alternative oxidase (AOX)	3V99 [22]	None	Dimer
Cytochrome P450 19A1 (CYP19A1)	3EQM [33]	2NZA [63]	Monomer
Fatty acid $\alpha$ -deoxygenase ( $\alpha$ -DOX)	4HHS [28]	1CXP [62]	Monomer
Fatty acid amide hydrolase (FAAH)	1MT5 [23]	1OCL [64]	Dimer
Flavoprotein-ubiquinone oxidoreductase (ETF-QO)	2GMH [65]	1PBE [66]	Monomer
Glycerol-3-phosphate dehydrogenase (GlpD)	2QCU [20]	4X9M [67]	Dimer
Inverting GT-B glycosyltransferase (WaaA)	2XCI [55]	1JG7 [68]	Monomer
Lysophosphatidic acid acyltransferase (PlsC)	5KYM [38]	1K30 [69]	Monomer
Phosphoglycosyl transferase (PglC)	5W7L [21]	None	Monomer
Squalene-hopene cyclase (SHC)	2SQC [31]	3DPY [70]	Dimer
Retaining GT-B glycosyltransferase (PglH)	6EJI [50]	1JG7 [68]	Dimer
Retaining GT-B glycosyltransferase (WbnH)	4XYW [48]	1JG7 [68]	Monomer
Retinoid isomerohydrolase P65 (RPE65)	4F2Z [24]	2BIW [25]	Dimer
Sulfide:quinone oxidoreductase (SQR)	3HYW [71]	1FCD [72]	Trimer
Tetraacyldisaccharide 4'-kinase (LpxK)	4EHW [54]	2HF9 [73]	Monomer
Type II NADH dehydrogenase (NDH-2)	4NWZ [74]	3GRS [75]	Dimer
Wall teichoic acid polymerase (TagF)	3L7I [76]	2IYA [77]	Dimer

The first 20 years of membrane protein structure determination yielded only four examples of monotopic folds, all of which act by extracting substrates from the membrane and extending membrane interactions by dimerization [1]. This initially gave a limited view of membrane-association strategies, which were shown later with additional examples to be more generalizable. The past decade has now yielded new examples that afford a greater diversity of both structure and the associated mechanisms of substrate interaction (see Figure 1C in Box 1).

### Something Old, Something New

A critical step in the biosynthesis and remodeling of phospholipids is catalyzed by the extensive family of 1-acyl-*sn*-glycerol-3-phosphate acyltransferases (Figure 3F). These monotopic enzymes simultaneously bind to the membrane-committed lysophospholipid substrate and the soluble acyl-thioester donor. Despite the importance of these enzymes, an experimentally determined structure has been elusive and attempts at structure prediction failed to yield a coherent solution that was consistent with a viable biochemical model [38]. The recent structure of PlsC from *Thermatoga maritima* now reveals that the functional domain of the enzyme is based on a known soluble  $\alpha\beta$ -acyltransferase fold with a signature HX<sub>4</sub>D motif, including the catalytic dyad, that is tightly associated with one face of the membrane by an intriguing two-helix motif. An extended N-terminal helix includes a core



## Trends in Biochemical Sciences

**Figure 3. Reactions of Structurally Characterized Monotopic Membrane Enzymes.** (A) Hydrolysis and alkene isomerization by RPE65; (B) COX-2, prostaglandin endoperoxide synthase; (C) α-DOX, linoleic acid α-hydroperoxidase; (D) SHC, squalene-hopene cyclase; (E) GlpD, FAD-dependent oxidation of glycerol 3-phosphate; (F) PlsC, lysophosphatidic acid acyltransferase; (G) steps in the *Campylobacter jejuni* protein glycosylation pathway – phosphoglycosyl transferase PglC and glycosyl transferases PglA, PglJ, and PglH; (H) steps in the *Escherichia coli* pathway for Lipid A biosynthesis, focusing on kinase LpxK and glycosyl transferase WaaA. Yellow highlighting in (G) and (H) emphasizes moieties with significant hydrophobic character.

of aromatic and other hydrophobic residues, delineated by basic residues, and is predicted to be positioned into a single leaflet of the bilayer, running parallel to the membrane interface (Figure 2). An adjacent partner helix is rich in basic residues, favoring placement at the membrane interface in the phospholipid head-group region. In contrast to the ‘extracting’ folds, a considerably higher percentage of the protein (16% calculated using the **PPM server**; Figure 1) is membrane-embedded. Thus, this intriguing, yet simple, architecture enables simultaneous interactions with both of the target substrates, despite their very different




properties, and sets the stage for ternary complex formation without requiring extraction of the amphipathic lysophospholipid from its membrane environment. The structure also provides insight into acyl chain specificity. The native specificity of PlsC is for a C16 palmitoyl group, which is shown to be perfectly accommodated in a hydrophobic pocket poised for transfer to the 2-position of the lysophospholipid substrate. The assignment of the acyl-binding site is corroborated by mutational studies that successfully switch the substrate specificity from palmitoyl to myristoyl transfer.

### A Marriage of Convenience

Monotopic membrane enzymes are prominent in biochemical pathways where localization of multiple enzymes, acting in sequence at the membrane interface, is highly advantageous. A common theme in the biogenesis of complex glycoconjugates is the stepwise assembly of glycans onto linear long-chain **polyprenol** phosphates [39]. For example, N-linked protein glycosylation in the *pgl* pathway of Gram-negative *Campylobacter* species involves the assembly of glycan-linked polyprenol diphosphate, through the action of the **phosphoglycosyl transferase (PGT)** PglC, followed by the **glycosyl transferases (GTases)** PglA, PglJ, PglH, and PglI on the cytoplasmic face of the inner bacterial membrane [40,41] (Figure 3G). These enzymes are predicted monotopic membrane enzymes that act directly and sequentially on amphiphilic polyprenol diphosphate-linked substrates, which remain resident in the membrane. The recently reported structure of PglC shows a novel membrane-association mode anchored by a **reentrant membrane helix** [15] that interacts with three short coplanar amphiphilic helices at the membrane interface [21]. The experimentally determined structure deviates from predictions based on primary structure and a hidden Markov model (**TMHMM**) [2], which suggested a bitopic architecture. PglC is the first characterized representative of a monotopic PGT superfamily and offers a unique structural view of a reentrant membrane helix supporting membrane association [21]. Based on extensive homology and covariance analysis, the reentrant membrane helix is conserved across this monotopic superfamily [42]. Structural and mechanistic studies on PglC revealed an active site, featuring an Asp–Glu dyad, perfectly positioned to capture the soluble uridine diphosphate bacillosamine (UDP-Bac) substrate for transfer of Bac-P to polyprenol phosphate (Pren-P) via a covalent enzyme intermediate [43] to afford Pren-PP-Bac. The structure of PglC shows no homology with known soluble folds or with the other superfamily of PGTs, which is exemplified by *MraY* and *WecA*, and features a polytopic architecture with 10 or 11 transmembrane helical domains (TMDs) and an active site crafted from inter-TMD loops [44,45].

Further assembly of the glycan-linked polyprenol diphosphate involves the action of a series of monotopic GTases based on a GT-B fold (Figure 3G) [46]. The GT-B-fold GTases are highly adaptable, and this soluble fold has evolved into peripheral, monotopic, and bitopic variants to meet functional requirements [47]. The GT-B fold displays C- and N-terminal **Rossmann-fold** domains, which meet at a central cleft containing the enzyme active site. The C-terminal domain binds the nucleoside diphosphate-sugar donor and the N-terminal domain, which associates with the membrane, binds the glycosyl moiety of the glycan-linked polyprenol diphosphate acceptor substrate. Thus, the hydrophobic polyprenol moiety positions the latter substrate for catalysis without the need for extraction from the membrane. Early in the *pgl* pathway, PglA transfers GalNAc to Pren-PP-Bac. Although PglA has yet to be crystallized, a functional homolog, *WbnH* (*Escherichia coli*), which catalyzes the transfer of GalNAc from UDP-GalNAc to Pren-PP-GalNAc to afford an  $\alpha$ -1,3-GalNAc linkage, has been characterized [48]. The membrane-associated structure of *WbnH*, predicted using the X-ray structure and the PPM web server [49], shows that *WbnH* associates with the membrane through three hydrophobic loops on the surface of the N-terminal domain (Figure 4A). This membrane-



## Trends in Biochemical Sciences

**Figure 4. Insight into GTase and Kinase Interactions with Membrane Calculated Using the PPM Server.** (A) GTase interaction with the membrane changes based on glycosyl acceptor substrate. Left: WbnH (4XYW) transfers GalNAc to membrane-resident Pren-PP-GalNAc. Right: PglH (6EJK) shown bound to nonhydrolyzable UDP-CH<sub>2</sub>-GalNAc (spheres) sequentially transfers three GalNAcs to membrane-resident Pren-PP-trisaccharide. C-terminal GT-B domain, pink; N-terminal GT-B domain, pale blue; bracing helix, pale green; membrane-interacting residues, slate blue; membrane surface, red; active-site cleft, red asterisk. (B) LpxK kinase changes conformation on binding to ADP-Mg<sup>2+</sup>. Protein colored N to C terminus, rainbow; membrane-associated residues, slate blue. Left: Open form of LpxK. Right: Closed form bound to ADP-Mg<sup>2+</sup>. ADP, space filling colored by atom; Mg<sup>2+</sup>, gray sphere. Red arrow shows movement of the C-terminal domain.

association domain is consistent with the C-terminal domain ‘dipping’ towards the membrane surface to accommodate the transfer of GalNAc to Pren-PP-monosaccharide at the active site situated between the protein domains.

PglH is a monotopic GT-B GTase that adds the fourth, fifth, and sixth sugars to the Pren-PP-trisaccharide (Figure 3G). PglH is intriguing as it adds three saccharide units in a processive fashion and reactions occur distal to the membrane surface. In this case, the predicted placement of PglH in the membrane calculated using the PPM server [49] reveals how the tilt of the structure relative to the membrane surface may accommodate these reactions by placing the active site cleft further from the membrane surface [50] compared with WbnH (Figure 4A). Insight into how PglH transfers predominantly three carbohydrate units comes from both mechanistic and structural analyses. Kinetic analysis with native substrates shows that each progressive sugar addition is retarded due to product inhibition, ultimately disfavoring further elongation of the product [51]. Additionally, recent structural analysis with substrates and substrate analogs suggests that specific interactions between the diphosphate moiety of the Pren-PP-glycan and basic residues on an  $\alpha$ -helix at the membrane surface act as a

molecular ruler that forms the basis for substrate specificity and determines the final product composition [50]. Ultimately, the placement of the extended polyprenol moiety of the substrate in the membrane, married to the distal active-site binding of the soluble glycan moiety, takes full advantage of the monotopic enzyme topology of PglH.

There are currently many aspects of monotopic GTase structure and function left to understand and a richer selection of structures in the fold family is needed. However, it is intriguing to speculate that the GT-B fold can be enlisted to transfer carbohydrates to specific positions in an elongating glycan chain based on the relative placement of the enzyme active site and the membrane surface.

### In the Neighborhood

Monotopic membrane enzymes also feature in early steps of the 'Raetz pathway' [52] for lipopolysaccharide (LPS) biosynthesis (Figure 3H). The first five steps of Kdo-Lipid A assembly occur in the cytosol. At this stage in the pathway, the intermediate has accumulated a preponderance of long-chain  $\beta$ -hydroxyacyl moieties and would not be free in solution; thus, the process transitions to a new neighborhood: the cytosolic face of the inner membrane. Tetraacyldisaccharide-1-phosphate 4'-kinase (LpxK) catalyzes ATP-dependent phosphorylation to afford the bisphosphorylated Lipid IV<sub>A</sub>. The structure of LpxK from *Aquifex aeolicus* (Figure 4B) reveals the first example of a P-loop kinase [53] adapted for function at the membrane interface through the addition of a membrane-associating N-terminal extension [54]. Thus, in this case, localization to the membrane occurs by sequence extension rather than by mutation in the fold, as in previously discussed examples. The *A. aeolicus* pathway is homologous to that from *E. coli*, showing only minor differences in the lengths of the  $\beta$ -hydroxyl-acyl tails installed [54]. The N-terminal helix of LpxK was originally predicted to be membrane spanning [2]; however, like PglC, interactions between the N terminus and the globular domain support a native monotopic topology and a functional monomer. Structural analysis of the ADP-Mg<sup>2+</sup>-bound enzyme shows engagement of the Walker A and B motif residues and supports a model wherein the tetraacylated disaccharide substrate is 'partially' extracted from the membrane to bring the C-4-hydroxyl group into proximity with the  $\gamma$ -phosphate of bound ATP (Figure 4B). Kinetic analysis supports the intermediacy of a ternary complex.

The LpxK product is elaborated by the monotopic GTase WaaA, which transfers a single Kdo (3-deoxy-D-manno-oct-2-ulosonic acid) from an activated CMP-Kdo donor. WaaA, like WbnH and PglH, is also a GT-B fold GTase. In this case, structural and computational studies illustrate how large-scale conformational changes may promote activity at the membrane interface without the need for extracting the amphiphilic substrate from its preferred environment: the N-terminal domain of WaaA associates with the membrane via a surface hydrophobic patch surrounded by a horseshoe-shaped ribbon of basic residues, which is proposed to facilitate access of the lipid A precursor to the active site at the interdomain cleft [55]. In WaaA, the cleft is large (23 Å × 20 Å) in the substrate-free state but closes down to position the catalytic machinery via the motion of two hinge regions (see Figure 1B in Box 1). Following WaaA are LpxL and LpxM, which are lauroyl and myristoyl acyltransferases belonging to a large superfamily of long-chain acyltransferases. The N-terminal hydrophobic domains of LpxL and LpxM had been previously predicted to form transmembrane helices, indicating a bitopic topology. Structures of full-length variants of this important superfamily have been difficult to obtain: the structure of the *Acinetobacter baumannii* LpxM was only recently reported [56]. The mode of membrane association for the fold cannot be inferred from the sole reported structure, in which the N-terminal domain protrudes from the soluble domain as a helix-break-helix motif.

However, recent biochemical and bioinformatics analyses of LpxM from *E. coli* now suggest a monotopic topology with the soluble domain anchored on the cytoplasmic face of the membrane by a reentrant helix [57].

### Concluding Remarks

Over the past decade the pace of membrane protein structure determination has increased due to collective advances in protein expression strategies, including cell-free methods, dedicated membrane protein crystallization screens, technical advances such as crystallization with endogenous lipids and in lipidic cubic phases, structural analysis in membrane-like scaffolds such as nanodiscs and bicelles, and a recent upsurge in the capabilities of, and access to, cryoelectron microscopy. However, despite these technological advances, structure determination of monotopic membrane proteins faces challenges specific to this class. In particular, creation of truncation constructs often utilizes natural domain boundaries to truncate proteins for improved purification and structure determination properties. However, these constructs often omit entirely or perturb membrane-associated domains. Therefore, while this strategy may lead to favorable crystallization and structure determination, omission of these domains leads to a loss of structural insight. The many examples described here, together with evolving methodologies in structural biology, now pave the way towards a deeper understanding of monotopic membrane proteins and their diversity of folds.

### Acknowledgments

Research related to monotopic membrane proteins in the Imperial and Allen groups has been supported by the NIH (GM-039334 to B.I.), the NIH Predoctoral Training Program (T32-GM007287 to S.E.), and the NIH Biomolecular Pharmacology Program Grant (T32-GM008541 to L.C.R.).

### References

1. Fomeris, F. and Mattevi, A. (2008) Enzymes without borders: mobilizing substrates, delivering products. *Science* 321, 213–216
2. Krogh, A. *et al.* (2001) Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J. Mol. Biol.* 305, 567–580
3. Almen, M.S. *et al.* (2009) Mapping the human membrane proteome: a majority of the human membrane proteins can be classified according to function and evolutionary origin. *BMC Biol.* 7, 50
4. Blobel, G. (1980) Intracellular protein topogenesis. *Proc. Natl. Acad. Sci. U. S. A.* 77, 1496–1500
5. Krebs, M.P. *et al.* (2004) Quality control of integral membrane proteins. *Trends Biochem. Sci.* 29, 648–655
6. Bracey, M.H. *et al.* (2004) Structural commonalities among integral membrane enzymes. *FEBS Lett.* 567, 159–165
7. Moravcevic, K. *et al.* (2012) Conditional peripheral membrane proteins: facing up to limited specificity. *Structure* 20, 15–27
8. Resh, M.D. (2013) Covalent lipid modifications of proteins. *Curr. Biol.* 23, R431–R435
9. Lomize, A.L. *et al.* (2007) The role of hydrophobic interactions in positioning of peripheral proteins in membranes. *BMC Struct. Biol.* 7, 44
10. Balali-Mood, K. *et al.* (2009) Interaction of monotopic membrane enzymes with a lipid bilayer: a coarse-grained MD simulation study. *Biochemistry* 48, 2135–2145
11. von Heijne, G. (2006) Membrane-protein topology. *Nat. Rev. Mol. Cell Biol.* 7, 909–918
12. Speers, A.E. and Wu, C.C. (2007) Proteomics of integral membrane proteins – theory and application. *Chem. Rev.* 107, 3687–3714
13. Fowler, P.W. *et al.* (2007) Monotopic enzymes and lipid bilayers: a comparative study. *Biochemistry* 46, 3108–3115
14. Yan, C. and Luo, J. (2010) An analysis of reentrant loops. *Protein J.* 29, 350–354
15. Viklund, H. *et al.* (2006) Structural classification and prediction of reentrant regions in  $\alpha$ -helical transmembrane proteins: application to complete genomes. *J. Mol. Biol.* 361, 591–603
16. Mattevi, A. (2006) A monotopic membrane protein goes solo. *Structure* 14, 628–629
17. Nam, H.J. *et al.* (2013) Rampant exchange of the structure and function of extramembrane domains between membrane and water soluble proteins. *PLoS Comput. Biol.* 9, e1002997
18. Nam, H.J. *et al.* (2015) Metazoans evolved by taking domains from soluble proteins to expand intercellular communication network. *Sci. Rep.* 5, 9576
19. Parsonage, D. *et al.* (1998) The soluble  $\alpha$ -glycerophosphate oxidase from *Enterococcus casseliflavus*. Sequence homology with the membrane-associated dehydrogenase and kinetic analysis of the recombinant enzyme. *J. Biol. Chem.* 273, 23812–23822
20. Yeh, J.I. *et al.* (2008) Structure of glycerol-3-phosphate dehydrogenase, an essential monotopic membrane enzyme involved in respiration and metabolism. *Proc. Natl. Acad. Sci. U. S. A.* 105, 3280–3285
21. Ray, L.C. *et al.* (2018) Membrane association of a monotopic phosphoglycosyl transferase underpins function. *Nat. Chem. Biol.* 4, 538–541
22. Shiba, T. *et al.* (2013) Structure of the trypanosome cyanide-insensitive alternative oxidase. *Proc. Natl. Acad. Sci. U. S. A.* 110, 4580–4585
23. Bracey, M.H. *et al.* (2002) Structural adaptations in a membrane enzyme that terminates endocannabinoid signaling. *Science* 298, 1793–1796
24. Kiser, P.D. *et al.* (2012) Structure of RPE65 isomerase in a lipidic matrix reveals roles for phospholipids and iron in catalysis. *Proc. Natl. Acad. Sci. U. S. A.* 109, E2747–E2756
25. Kloer, D.P. *et al.* (2005) The structure of a retinal-forming carotenoid oxygenase. *Science* 308, 267–269

### Outstanding Questions

What mechanisms might allow the evolution of monotopic membrane proteins besides the mere recruitment of soluble folds to the membrane?

How does the evolution of shallowly embedded monotopic proteins differ from that of proteins that penetrate more deeply into the hydrophobic core of the membrane?

Why have monotopic membrane proteins with reentrant helical domains proved to be so intractable to structure determination in the past? How can recent advances in the field be applied to expand the diversity of monotopic membrane proteins in the PDB?

How can recent advances in techniques such as cryoelectron microscopy and the use of model membrane scaffolds, such as nanodiscs and lipidic cubic phases, be leveraged to yield the most relevant structures of membrane-embedded proteins in native-like environments?

Are there specific motifs driving monotopic membrane association via reentrant helices? Can the knowledge of these motifs be leveraged to predict membrane association modes for structurally and functionally uncharacterized proteins?

How does the composition of the lipid bilayer, with respect to the phospholipid head groups, acyl chains, and lipid additives, affect enzyme activity?

Can the current understanding of the function of monotopic enzymes inform bioengineering efforts to apply enzymes as catalysts for transformations on hydrophobic or amphiphilic substrates?

26. Picot, D. *et al.* (1994) The X-ray crystal structure of the membrane protein prostaglandin H2 synthase-1. *Nature* 367, 243–249
27. Zeng, J. and Fenna, R.E. (1992) X-ray crystal structure of canine myeloperoxidase at 3 Å resolution. *J. Mol. Biol.* 226, 185–207
28. Goulah, C.C. *et al.* (2013) The crystal structure of  $\alpha$ -dioxygenase provides insight into diversity in the cyclooxygenase-peroxidase superfamily. *Biochemistry* 52, 1364–1372
29. Sanz, A. *et al.* (1998) PLOX, a new pathogen-induced oxygenase with homology to animal cyclooxygenase. *Plant Cell* 10, 1523–1537
30. Wendt, K.U. *et al.* (1997) Structure and function of a squalene cyclase. *Science* 277, 1811–1815
31. Wendt, K.U. *et al.* (1999) The structure of the membrane protein squalene-hopene cyclase at 2.0 Å resolution. *J. Mol. Biol.* 286, 175–187
32. Thoma, R. *et al.* (2004) Insight into steroid scaffold formation from the structure of human oxidosqualene cyclase. *Nature* 432, 118–122
33. Ghosh, D. *et al.* (2009) Structural basis for androgen specificity and oestrogen synthesis in human aromatase. *Nature* 457, 219–223
34. Denisov, I.G. *et al.* (2012) Structural differences between soluble and membrane bound cytochrome P450s. *J. Inorg. Biochem.* 108, 150–158
35. Johnson, E.F. and Stout, C.D. (2013) Structural diversity of eukaryotic membrane cytochrome P450s. *J. Biol. Chem.* 288, 17082–17090
36. Cojocaru, V. *et al.* (2011) Structure and dynamics of the membrane-bound cytochrome P450 2C9. *PLoS Comput. Biol.* 7, e1002152
37. Monk, B.C. *et al.* (2014) Architecture of a single membrane spanning cytochrome P450 suggests constraints that orient the catalytic domain relative to a bilayer. *Proc. Natl. Acad. Sci. U. S. A.* 111, 3865–3870
38. Robertson, R.M. *et al.* (2017) A two-helix motif positions the lysophosphatidic acid acyltransferase active site for catalysis within the membrane bilayer. *Nat. Struct. Mol. Biol.* 24, 666–671
39. Hartley, M.D. and Imperiali, B. (2012) At the membrane frontier: a prospectus on the remarkable evolutionary conservation of polyprenols and polyprenyl-phosphates. *Arch. Biochem. Biophys.* 517, 83–97
40. Szymanski, C.M. *et al.* (1999) Evidence for a system of general protein glycosylation in *Campylobacter jejuni*. *Mol. Microbiol.* 32, 1022–1030
41. Glover, K.J. *et al.* (2005) *In vitro* assembly of the undecaprenylpyrophosphate-linked heptasaccharide for prokaryotic N-linked glycosylation. *Proc. Natl. Acad. Sci. U. S. A.* 102, 14255–14259
42. Lukose, V. *et al.* (2015) Conservation and covariance in small bacterial phosphoglycosyltransferases identify the functional catalytic core. *Biochemistry* 54, 7326–7334
43. Das, D. *et al.* (2017) Analysis of a dual domain phosphoglycosyl transferase reveals a ping-pong mechanism with a covalent enzyme intermediate. *Proc. Natl. Acad. Sci. U. S. A.* 114, 7019–7024
44. Chung, B.C. *et al.* (2013) Crystal structure of MraY, an essential membrane enzyme for bacterial cell wall synthesis. *Science* 341, 1012–1016
45. Lukose, V. *et al.* (2017) Bacterial phosphoglycosyl transferases: initiators of glycan biosynthesis at the membrane interface. *Glycobiology* 27, 820–833
46. Lairson, L.L. *et al.* (2008) Glycosyltransferases: structures, functions, and mechanisms. *Annu. Rev. Biochem.* 77, 521–555
47. Albesa-Jove, D. *et al.* (2014) Structure–function relationships of membrane-associated GT-B glycosyltransferases. *Glycobiology* 24, 108–124
48. Li, F. *et al.* (2015) The structure of WbnH in a near active state. *Protein Cell* 6, 615–618
49. Lomize, M.A. *et al.* (2012) OPM database and PPM web server: resources for positioning of proteins in membranes. *Nucleic Acids Res.* 40, D370–D376
50. Ramirez, A.S. *et al.* (2018) Structural basis of the molecular ruler mechanism of a bacterial glycosyltransferase. *Nat. Commun.* 9, 445
51. Troutman, J.M. and Imperiali, B. (2009) *Campylobacter jejuni* PglH is a single active site processive polymerase that utilizes product inhibition to limit sequential glycosyl transfer reactions. *Biochemistry* 48, 2807–2816
52. Raetz, C.R. and Whitfield, C. (2002) Lipopolysaccharide endotoxins. *Annu. Rev. Biochem.* 71, 635–700
53. Leipe, D.D. *et al.* (2003) Evolution and classification of P-loop kinases and related proteins. *J. Mol. Biol.* 333, 781–815
54. Emptage, R.P. *et al.* (2012) Crystal structure of LpxK, the 4'-kinase of Lipid A biosynthesis and atypical P-loop kinase functioning at the membrane interface. *Proc. Natl. Acad. Sci. U. S. A.* 109, 12956–12961
55. Schmidt, H. *et al.* (2012) Structural and mechanistic analysis of the membrane-embedded glycosyltransferase WaaA required for lipopolysaccharide synthesis. *Proc. Natl. Acad. Sci. U. S. A.* 109, 6253–6258
56. Dovala, D. *et al.* (2016) Structure-guided enzymology of the lipid A acyltransferase LpxM reveals a dual activity mechanism. *Proc. Natl. Acad. Sci. U. S. A.* 113, E6064–E6071
57. Entova, S. *et al.* (2018) Insights into the key determinants of membrane protein topology enable the identification of new monotopic folds. *Elife* 7, e40889
58. Magotti, P. *et al.* (2015) Structure of human N-acylphosphatidylethanolamine-hydrolyzing phospholipase D: regulation of fatty acid ethanolamide biosynthesis by bile acids. *Structure* 23, 598–604
59. Carfi, A. *et al.* (1995) The 3-D structure of a zinc metallo-beta-lactamase from *Bacillus cereus* reveals a new type of protein fold. *EMBO J.* 14, 4914–4921
60. Hsiao, Y.S. *et al.* (2006) Crystal structure of rat carnitine palmitoyltransferase II (CPT-II). *Biochem. Biophys. Res. Commun.* 346, 974–980
61. Jogl, G. and Tong, L. (2003) Crystal structure of carnitine acetyltransferase and implications for the catalytic mechanism and fatty acid transport. *Cell* 112, 113–122
62. Fiedler, T.J. *et al.* (2000) X-ray crystal structure and characterization of halide-binding sites of human myeloperoxidase at 1.8 Å resolution. *J. Biol. Chem.* 275, 11964–11971
63. Zhao, B. *et al.* (2007) Different binding modes of two flavin substrate molecules in cytochrome P450 158A1 (CYP158A1) compared to CYP158A2. *Biochemistry* 46, 8725–8733
64. Shin, S. *et al.* (2003) Characterization of a novel Ser–cisSer–Lys catalytic triad in comparison with the classical Ser–His–Asp triad. *J. Biol. Chem.* 278, 24937–24943
65. Zhang, J. *et al.* (2006) Structure of electron transfer flavoprotein-ubiquinone oxidoreductase and electron transfer to the mitochondrial ubiquinone pool. *Proc. Natl. Acad. Sci. U. S. A.* 103, 16212–16217
66. Schreuder, H.A. *et al.* (1989) Crystal structure of the *p*-hydroxybenzoate hydroxylase-substrate complex refined at 1.9 Å resolution. Analysis of the enzyme–substrate and enzyme–product complexes. *J. Mol. Biol.* 208, 679–696
67. Elkhail, C.K. *et al.* (2015) Structure and proposed mechanism of L- $\alpha$ -glycerophosphate oxidase from *Mycoplasma pneumoniae*. *FEBS J.* 282, 3030–3042
68. Morera, S. *et al.* (2001) High resolution crystal structures of T4 phage  $\beta$ -glucosyltransferase: induced fit and effect of substrate and metal binding. *J. Mol. Biol.* 311, 569–577
69. Turnbull, A.P. *et al.* (2001) Analysis of the structure, substrate specificity, and mechanism of squash glycerol-3-phosphate (1)-acyltransferase. *Structure* 9, 347–353
70. DeGraw, A.J. *et al.* (2008) Caged protein prenyltransferase substrates: tools for understanding protein prenylation. *Chem. Biol. Drug Des.* 72, 171–181

71. Marcia, M. *et al.* (2009) The structure of *Aquifex aeolicus* sulfide: quinone oxidoreductase, a basis to understand sulfide detoxification and respiration. *Proc. Natl. Acad. Sci. U. S. A.* 106, 9625–9630
72. Chen, Z.W. *et al.* (1994) The structure of flavocytochrome c sulfide dehydrogenase from a purple phototrophic bacterium. *Science* 266, 430–432
73. Gasper, R. *et al.* (2006) Structural insights into HypB, a GTP-binding protein that regulates metal binding. *J. Biol. Chem.* 281, 27492–27502
74. Heikal, A. *et al.* (2014) Structure of the bacterial type II NADH dehydrogenase: a monotopic membrane protein with an essential role in energy generation. *Mol. Microbiol.* 91, 950–964
75. Karplus, P.A. and Schulz, G.E. (1987) Refined structure of glutathione reductase at 1.54 Å resolution. *J. Mol. Biol.* 195, 701–729
76. Lovering, A.L. *et al.* (2010) Structure of the bacterial teichoic acid polymerase TagF provides insights into membrane association and catalysis. *Nat. Struct. Mol. Biol.* 17, 582–589
77. Bolam, D.N. *et al.* (2007) The crystal structure of two macrolide glycosyltransferases provides a blueprint for host cell antibiotic immunity. *Proc. Natl. Acad. Sci. U. S. A.* 104, 5336–5341